

AP[®] CHEMISTRY
2007 SCORING GUIDELINES (Form B)

Question 1 (continued)

- (d) State whether the entropy change, ΔS° , for the reaction at 862°C is positive, negative, or zero. Justify your answer.

ΔS° is <u>positive</u> because four moles of gaseous products are produced from three moles of gaseous reactants.	One point is earned for the correct explanation.
--	--

- (e) State whether the enthalpy change, ΔH° , for the reaction at 862°C is positive, negative, or zero. Justify your answer.

Both ΔG° and ΔS° are positive, as determined in parts (c) and (d). Thus, ΔH° must be positive because ΔH° is the sum of two positive terms in the equation $\Delta H^\circ = \Delta G^\circ + T\Delta S^\circ$.	One point is earned for the correct sign. One point is earned for a correct explanation.
--	---

- (f) After a certain period of time, 1.000 mol of $O_2(g)$ is added to the mixture in the flask. Does the mass of $U_3O_8(s)$ in the flask increase, decrease, or remain the same? Justify your answer.

The mass of $U_3O_8(s)$ will <u>increase</u> because the reaction is at equilibrium, and the addition of a product creates a “stress” on the product (right) side of the reaction. The reaction will then proceed from right to left to reestablish equilibrium so that some $O_2(g)$ is consumed (tending to relieve the stress) as more $U_3O_8(s)$ is produced.	One point is earned for a correct explanation.
--	--

1A,

CHEMISTRY
Section II
(Total time—95 minutes)


Part A
Time—55 minutes
YOU MAY USE YOUR CALCULATOR FOR PART A.

CLEARLY SHOW THE METHOD USED AND THE STEPS INVOLVED IN ARRIVING AT YOUR ANSWERS. It is to your advantage to do this, since you may obtain partial credit if you do and you will receive little or no credit if you do not. Attention should be paid to significant figures.

Be sure to write all your answers to the questions on the lined pages following each question in this booklet. Do NOT write your answers on the lavender insert.

Answer Questions 1, 2, and 3. The Section II score weighting for each question is 20 percent.

1. A sample of solid U_3O_8 is placed in a rigid 1.500 L flask. Chlorine gas, $\text{Cl}_2(\text{g})$, is added, and the flask is heated to 862°C . The equation for the reaction that takes place and the equilibrium-constant expression for the reaction are given below.



When the system is at equilibrium, the partial pressure of $\text{Cl}_2(\text{g})$ is 1.007 atm and the partial pressure of $\text{UO}_2\text{Cl}_2(\text{g})$ is 9.734×10^{-4} atm.

- (a) Calculate the partial pressure of $\text{O}_2(\text{g})$ at equilibrium at 862°C .
- (b) Calculate the value of the equilibrium constant, K_p , for the system at 862°C .
- (c) Calculate the Gibbs free-energy change, ΔG° , for the reaction at 862°C .
- (d) State whether the entropy change, ΔS° , for the reaction at 862°C is positive, negative, or zero. Justify your answer.
- (e) State whether the enthalpy change, ΔH° , for the reaction at 862°C is positive, negative, or zero. Justify your answer.
- (f) After a certain period of time, 1.000 mol of $\text{O}_2(\text{g})$ is added to the mixture in the flask. Does the mass of $\text{U}_3\text{O}_8(\text{s})$ in the flask increase, decrease, or remain the same? Justify your answer.

a) partial pressure is proportional to number of mol

3 mol UO_2Cl_2 : 1 mol O_2

$$\frac{1}{3} (9.734 \times 10^{-4} \text{ atm}) = 3.245 \times 10^{-4} \text{ atm}$$

$$(b) K_p = \frac{(9.734 \times 10^{-4})^3 (3.245 \times 10^{-4})}{(1.007)^3}$$

$$= 2.931 \times 10^{-13}$$

(c) at equilibrium, $\Delta G^\circ = -RT \ln K$

$$R = 0.00831 \text{ kJ mol}^{-1} \text{ K}^{-1}$$

$$T = 1135 \text{ K}$$

$$K = 2.931 \times 10^{-13}$$

$$\Delta G^\circ = -(0.00831)(1135) \ln(2.931 \times 10^{-13})$$

$$= 272.2 \text{ kJ/mol}$$

(d) Positive; a solid and a gas form two gases, 3 mol of gas forms 4 mol of gas.

$$(e) \Delta G^\circ = \Delta H^\circ - T \Delta S^\circ$$

ΔG° is positive

ΔS° is positive

T is positive

$$(+)=\Delta H^\circ-(+)$$

Positive; in order for the equation to remain true.

(f) Increase; adding more $O_2(g)$ shifts the equilibrium to the left, producing more mol $H_2O(g)$ and thus a greater mass as well.

181

CHEMISTRY
Section II
(Total time—95 minutes)


Part A
Time—55 minutes
YOU MAY USE YOUR CALCULATOR FOR PART A.

CLEARLY SHOW THE METHOD USED AND THE STEPS INVOLVED IN ARRIVING AT YOUR ANSWERS. It is to your advantage to do this, since you may obtain partial credit if you do and you will receive little or no credit if you do not. Attention should be paid to significant figures.

Be sure to write all your answers to the questions on the lined pages following each question in this booklet. Do NOT write your answers on the lavender insert.

Answer Questions 1, 2, and 3. The Section II score weighting for each question is 20 percent.

1. A sample of solid U_3O_8 is placed in a rigid, 1.500 L flask. Chlorine gas, $Cl_2(g)$, is added, and the flask is heated to $862^\circ C$. The equation for the reaction that takes place and the equilibrium-constant expression for the reaction are given below.



When the system is at equilibrium, the partial pressure of $Cl_2(g)$ is 1.007 atm and the partial pressure of $UO_2Cl_2(g)$ is 9.734×10^{-4} atm.

- (a) Calculate the partial pressure of $O_2(g)$ at equilibrium at $862^\circ C$
- (b) Calculate the value of the equilibrium constant, K_p , for the system at $862^\circ C$.
- (c) Calculate the Gibbs free-energy change, ΔG° , for the reaction at $862^\circ C$.
- (d) State whether the entropy change, ΔS° , for the reaction at $862^\circ C$ is positive, negative, or zero. Justify your answer.
- (e) State whether the enthalpy change, ΔH° , for the reaction at $862^\circ C$ is positive, negative, or zero. Justify your answer.
- (f) After a certain period of time, 1.000 mol of $O_2(g)$ is added to the mixture in the flask. Does the mass of $U_3O_8(s)$ in the flask increase, decrease, or remain the same? Justify your answer.

a) $PV = nRT$ $P = \frac{nRT}{V}$ $1 \times 0.0821 \times 1135 = 62.1 \text{ atm}$

$$b) K_p = \frac{(9.734 \times 10^{-4})^3 (62.1)}{(1.007)^3} = 5.609 \times 10^{-8}$$

$$c) \Delta G^\circ = -RT \ln K$$

$$= -8.314 \text{ J mol}^{-1} \text{ K}^{-1} \times 1135 \times \ln(5.609 \times 10^{-8})$$

$$= 157552.8 \text{ J} \div 1000$$

$$= +157.55 \text{ kJ}$$

d) positive

There is a phase change when U_3O_8 (solid) reacts with Cl_2 : It becomes gas. As solid becomes gas, the entropy increases as the randomness of gas is greater than that of the solid.

$$e) \Delta G^\circ = \Delta H^\circ - T \Delta S^\circ$$

$$175.55 = \Delta H^\circ - 1135 \Delta S^\circ$$

$$\Delta H^\circ = \text{positive}$$

ΔH° is positive because when U_3O_8 reacts with Cl_2 , it requires energy and the heat is absorbed = endothermic

f) The mass of U_3O_8 stays the same. This is because 1.00 mol of O_2 is negligibly small so it does not affect the mass of U_3O_8

$$K_p = 9.05 \times 10^{-10}$$

$$\begin{aligned} \Delta G &= -RT \ln K \\ &= -2.303 RT \log K \end{aligned}$$

$$\begin{aligned} \Delta G &= -2.303 \cdot .0821 \times (862 + 273K) \times \log (9.05 \times 10^{-10}) \\ \Delta G &= +1940 \text{ kJ} \end{aligned}$$

d) It is negative because $\Delta G = \Delta H - T\Delta S$ means that ΔS has to be a negative for the ΔG to be positive as it is.

e) Negative as the reaction is exothermic and releases heat.

f) Decreases because it is in equilibrium and any extra mass takes up its space in the reaction, since it is added to the same side.

AP[®] CHEMISTRY
2007 SCORING COMMENTARY (Form B)

Question 1

Sample: 1A

Score: 9

This response earned all 9 points: 1 for part (a), 2 for part (b), 2 for part (c), 1 for part (d), 2 for part (e), and 1 for part (f).

Sample: 1B

Score: 6

The point was not earned in part (a) because the student incorrectly attempts to use the Ideal Gas Law to determine the partial pressure of O₂ gas at equilibrium. Both points were earned in part (b) because the student correctly substitutes and calculates a value for the equilibrium constant using the pressure calculated in part (a). Both points were earned in part (c) because the student correctly substitutes and calculates a value for ΔG° using the K_p from part (b). The point was earned in part (d). The first point was earned for part (e) because the student correctly identifies the sign of ΔH° as positive; however, the second point was not earned because the justification given relates to the heat added to start the reaction rather than a connection to the Gibb's equation. The point was not earned in part (f).

Sample: 1C

Score: 3

The point was not earned in part (a). Both points were earned in part (b) because the student correctly substitutes and calculates a value for the equilibrium constant using the calculated pressure from part (a). Only 1 point was earned in part (c) because although the student substitutes correctly, using the K_p value calculated in part (b), the incorrect R is used in the equation. No points were earned for parts (d), (e), or (f).