

2022

AP[®]

 CollegeBoard

AP[®] Biology

Free-Response Questions

© 2022 College Board. College Board, Advanced Placement, AP, AP Central, and the acorn logo are registered trademarks of College Board. Visit College Board on the web: collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

AP® BIOLOGY EQUATIONS AND FORMULAS

Statistical Analysis and Probability

Mean

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

Standard Deviation

$$s = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n - 1}}$$

Standard Error of the Mean

$$SE_{\bar{x}} = \frac{s}{\sqrt{n}}$$

Chi-Square

$$\chi^2 = \sum \frac{(o - e)^2}{e}$$

Chi-Square Table

p value	Degrees of Freedom							
	1	2	3	4	5	6	7	8
0.05	3.84	5.99	7.81	9.49	11.07	12.59	14.07	15.51
0.01	6.63	9.21	11.34	13.28	15.09	16.81	18.48	20.09

Laws of Probability

If A and B are mutually exclusive, then:

$$P(A \text{ or } B) = P(A) + P(B)$$

If A and B are independent, then:

$$P(A \text{ and } B) = P(A) \times P(B)$$

Hardy-Weinberg Equations

$$p^2 + 2pq + q^2 = 1 \quad p = \text{frequency of allele 1 in a population}$$

$$p + q = 1 \quad q = \text{frequency of allele 2 in a population}$$

\bar{x} = sample mean

n = sample size

s = sample standard deviation (i.e., the sample-based estimate of the standard deviation of the population)

o = observed results

e = expected results

Σ = sum of all

Degrees of freedom are equal to the number of distinct possible outcomes minus one.

Metric Prefixes

<u>Factor</u>	<u>Prefix</u>	<u>Symbol</u>
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^{-1}	deci	d
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Mode = value that occurs most frequently in a data set

Median = middle value that separates the greater and lesser halves of a data set

Mean = sum of all data points divided by number of data points

Range = value obtained by subtracting the smallest observation (sample minimum) from the greatest (sample maximum)

Rate and Growth		Water Potential (Ψ)
<p>Rate</p> $\frac{dY}{dt}$ <p>Population Growth</p> $\frac{dN}{dt} = B - D$ <p>Exponential Growth</p> $\frac{dN}{dt} = r_{\max} N$ <p>Logistic Growth</p> $\frac{dN}{dt} = r_{\max} N \left(\frac{K - N}{K} \right)$	<p>dY = amount of change</p> <p>dt = change in time</p> <p>B = birth rate</p> <p>D = death rate</p> <p>N = population size</p> <p>K = carrying capacity</p> <p>r_{\max} = maximum per capita growth rate of population</p>	<p>$\Psi = \Psi_P + \Psi_S$</p> <p>Ψ_P = pressure potential</p> <p>Ψ_S = solute potential</p> <p>The water potential will be equal to the solute potential of a solution in an open container because the pressure potential of the solution in an open container is zero.</p> <p>The Solute Potential of a Solution</p> <p>$\Psi_S = -iCRT$</p> <p>i = ionization constant (1.0 for sucrose because sucrose does not ionize in water)</p> <p>C = molar concentration</p> <p>R = pressure constant ($R = 0.0831$ liter bars/mole K)</p> <p>T = temperature in Kelvin ($^{\circ}\text{C} + 273$)</p>
<p>Simpson's Diversity Index</p> <p>Diversity Index = $1 - \sum \left(\frac{n}{N} \right)^2$</p> <p>$n$ = total number of organisms of a particular species</p> <p>N = total number of organisms of all species</p>		<p>pH = $-\log[\text{H}^+]$</p>
Surface Area and Volume		
<p>Surface Area of a Sphere</p> $SA = 4\pi r^2$ <p>Surface Area of a Rectangular Solid</p> $SA = 2lh + 2lw + 2wh$ <p>Surface Area of a Cylinder</p> $SA = 2\pi rh + 2\pi r^2$ <p>Surface Area of a Cube</p> $SA = 6s^2$	<p>Volume of a Sphere</p> $V = \frac{4}{3}\pi r^3$ <p>Volume of a Rectangular Solid</p> $V = lwh$ <p>Volume of a Cylinder</p> $V = \pi r^2 h$ <p>Volume of a Cube</p> $V = s^3$	<p>r = radius</p> <p>l = length</p> <p>h = height</p> <p>w = width</p> <p>s = length of one side of a cube</p> <p>SA = surface area</p> <p>V = volume</p>

BIOLOGY

SECTION II

Time—1 hour and 30 minutes

6 Questions

Directions: Questions 1 and 2 are long free-response questions that require about 25 minutes each to answer. Questions 3 through 6 are short free-response questions that require about 10 minutes each to answer.

Read each question carefully and completely. Answers must be written out in paragraph form. Outlines, bulleted lists, or diagrams alone are not acceptable.

You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

Question 1 is on the following page.

1. The binding of an extracellular ligand to a G protein-coupled receptor in the plasma membrane of a cell triggers intracellular signaling (Figure 1, A). After ligand binding, GTP replaces the GDP that is bound to $G\alpha$, a subunit of the G protein (Figure 1, B). This causes $G\alpha$ to activate other cellular proteins, including adenylyl cyclase that converts ATP to cyclic AMP (cAMP). The cAMP activates protein kinases (Figure 1, C). In cells that line the small intestine, a cAMP-activated protein kinase causes further signaling that ultimately results in the secretion of chloride ions (Cl^-) from the cells. Under normal conditions, $G\alpha$ hydrolyzes GTP to GDP, thus inactivating adenylyl cyclase and stopping the signal (Figure 1, A).




Figure 1. Under normal conditions, ligand binding to a G protein-coupled receptor results in chloride ion transport from an intestinal cell.

Individuals infected with the bacterium *Vibrio cholerae* experience severe loss of water from the body (dehydration). This is due to the effects of the bacterial cholera toxin that enters intestinal cells. Scientists studied the effects of cholera toxin on four samples of isolated intestinal cell membranes containing the G protein-related signal transduction components shown in Figure 1. GTP was added to samples II and IV only; cholera toxin was added to samples III and IV only. The scientists then measured the amount of cAMP produced by the adenylyl cyclase in each sample (Table 1).

TABLE 1. AMOUNT OF cAMP PRODUCED FROM INTESTINAL CELL MEMBRANES IN THE ABSENCE OR PRESENCE OF CHOLERA TOXIN

Sample	GTP	Cholera Toxin	Rate of cAMP Production (pmol per mg adenylyl cyclase per min)
I	–	–	0.5
II	+	–	10.0
III	–	+	0.5
IV	+	+	127.0

present, +; absent, –

- (a) **Describe** one characteristic of a membrane that requires a channel be present for chloride ions to passively cross the membrane. **Explain** why the movement of chloride ions out of intestinal cells leads to water loss.
- (b) **Identify** an independent variable in the experiment. **Identify** a negative control in the experiment. **Justify** why the scientists included Sample III as a control treatment in the experiment.
- (c) Based on the data, **describe** the effect of cholera toxin on the synthesis of cAMP. **Calculate** the percent change in the rate of cAMP production due to the presence of cholera toxin in sample IV compared with sample II.
- (d) A drug is designed to bind to cholera toxin before it crosses the intestinal cell membrane. Scientists mix the drug with cholera toxin and then add this mixture and GTP to a sample of intestinal cell membranes. **Predict** the rate of cAMP production in pmol per mg adenylyl cyclase per min if the drug binds to all of the toxin. In a separate experiment, scientists engineer a mutant adenylyl cyclase that cannot be activated by $Gs\alpha$. The scientists claim that cholera toxin will not cause excessive water loss from whole intestinal cells that contain the mutant adenylyl cyclase. **Justify** this claim.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

2. During meiosis, double-strand breaks occur in chromatids. The breaks are either repaired by the exchange of genetic material between homologous nonsister chromatids, which is the process known as crossing over (Figure 1A), or they are simply repaired without any crossing over (Figure 1B). Plant breeders developing new varieties of corn are interested in determining whether, in corn, a correlation exists between the number of meiotic double-strand chromatid breaks and the number of crossovers.




Figure 1. Double-strand breaks in chromatids are repaired with crossing over (A) or without crossing over (B).

Using specialized staining and microscopy techniques, scientists counted the number of double-strand chromatid breaks and the number of crossovers in the same number of meiotic gamete-forming cells of six inbred strains of corn (Table 1).

TABLE 1. NUMBER OF CHROMATID DOUBLE-STRAND BREAKS AND AVERAGE NUMBER OF CROSSOVERS IN INBRED STRAINS OF CORN

Strain of Corn	Number of Double-Strand Breaks	Average Number of Crossovers ($\pm 2SE_{\bar{x}}$)
I	710	19.5 ± 0.5
II	650	18.0 ± 0.7
III	600	17.5 ± 1.0
IV	510	16.0 ± 1.0
V	425	14.0 ± 0.5
VI	325	11.0 ± 1.5

- (a) The double-strand breaks occur along the DNA backbone. **Describe** the process by which the breaks occur.
- (b) Using the template in the space provided for your response, **construct** an appropriately labeled graph that represents the data in Table 1 and allows examination of a possible correlation between double-strand breaks and crossovers. Based on the data, **determine** whether corn strains I, II, and III differ in their average number of crossovers.
- (c) Based on the data, **describe** the relationship between the average number of double-strand breaks and the average number of crossovers in the strains of corn analyzed in the experiment.
- (d) Crossing over (Figure 1A) creates physical connections that are required for proper separation of homologous chromosomes during meiosis. A diploid cell with four pairs of homologous chromosomes undergoes meiosis to produce four haploid cells. Crossing over occurs between only three of the pairs. **Predict** the number of chromosomes most likely present in each of the four haploid cells. Provide reasoning to **justify** your prediction. **Explain** how plant breeders can use the information in Table 1 to help develop new varieties of corn.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

3. Fireflies emit light when the enzyme luciferase catalyzes a reaction in which its substrate, D-luciferin, reacts to form oxyluciferin and other products (Figure 1). In order to determine the optimal temperature for this enzyme, scientists added ATP to a solution containing D-luciferin, luciferase, and other substances needed for the reaction. They then measured the amount of light emitted during the first three seconds of the reaction when it was carried out at different temperatures.




Figure 1. Light is emitted as a result of the reaction catalyzed by luciferase.

- (a) **Describe** a characteristic of the luciferase enzyme that allows it to catalyze the reaction.
- (b) **Identify** the dependent variable in the experiment.
- (c) **State** the null hypothesis for the experiment.
- (d) A student claims that, as temperature increases, there will be an increase in the amount of light given off by the reaction in the first three seconds. **Support** the student's claim.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

4. Existing isolated brook trout populations in Newfoundland, Canada, were once part of a larger population that was fragmented at the end of the most recent glaciation period about 10,000 to 12,000 years ago. Researchers investigated 14 naturally separated stream populations of brook trout. They found that the populations are all genetically distinct and show differences in morphology.
- (a) **Describe** the prezygotic barrier that results in these genetically distinct populations.
 - (b) Brook trout with longer fins are able to swim faster than brook trout with shorter fins. In one of the Newfoundland streams, the main prey of the brook trout evolved to move faster. For brook trout living in this stream, **explain** the difference in fitness between longer-finned individuals and shorter-finned individuals.
 - (c) If two morphologically and behaviorally distinct populations of brook trout remain isolated for many generations, **predict** the likely impact on both populations.
 - (d) Researchers claim that there are more genetic differences between any two current brook trout populations than there are between any single current population and the ancestral brook trout population from which all the trout are descended. Provide reasoning to **justify** their claim.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

5. The following models represent all the interacting species in two different communities with some of the same species and feeding relationships. These models assume that both communities have the same initial biomass. The models can be used to understand the effects of human activities on the communities.




Figure 1. Models of two different communities with some of the same species

- (a) **Describe** a characteristic of a community that makes a species invasive in that community but not invasive in a different community.
- (b) **Explain** why removing species PP1 will have a greater effect on community B than on community A.
- (c) An invasive species (INV) that eats individuals of species SC2 is introduced into community B. Using the template in the space provided for your response, for community B, indicate the feeding relationship for this invasive species by correctly placing **INV** to represent the invasive species and an **arrow** to represent the feeding relationship within community B.
- (d) **Explain** how human activities that add toxins to the soil could change a community with many species at each trophic level, such as community A, into a community with few species at each trophic level, such as community B.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

6. Researchers are studying the use of RNA vaccines to protect individuals against certain diseases. To develop the vaccines, particular cells are first removed from an individual. Then mRNAs coding for specific proteins from a pathogen are introduced into the cells. The altered cells are injected back into the individual, where the cells make the proteins encoded by the introduced mRNAs. The individual then produces an immune response to the proteins that will help to protect the individual from developing a disease if exposed to the pathogen in the future.

When introduced into cells, the mRNAs used for vaccines must be stable so that they are not degraded before the encoded proteins are produced. Researchers developed several modified caps that they hypothesized might make the introduced mRNAs more stable than mRNAs with the normal GTP cap. To test the effect of the modified caps, the researchers produced mRNAs that differed only in their cap structure (no cap, the normal cap, or modified caps I, II, or III). They introduced the same amount of each mRNA to different groups of cells and measured the amount of time required for half of the mRNAs to degrade (mRNA half-life) and the total amount of protein translated from the mRNAs (Table 1).

TABLE 1. EFFECT OF mRNA CAP STRUCTURE ON mRNA HALF-LIFE AND PROTEIN TRANSLATED FROM THE INTRODUCED mRNA

5' Cap Structure	mRNA Half-Life $\pm 2SE_{\bar{x}}$ (hours after introduction into cells)	Total Amount of Protein Translated from mRNA $\pm 2SE_{\bar{x}}$ (relative to amount in normal cap)
No cap	1.41 ± 0.02	0.011 ± 0.000
Normal GTP cap	16.10 ± 1.83	1.000 ± 0.007
Modified cap I	15.50 ± 1.57	4.777 ± 0.042
Modified cap II	27.00 ± 2.85	13.094 ± 0.307
Modified cap III	18.09 ± 0.81	6.570 ± 0.075

- (a) Based on the data, **identify** which cap structure is most likely to protect the end of the mRNAs from degradation.
- (b) Based on the data for the mRNAs with modified caps, **describe** the relationship between the mRNA half-life and the total amount of protein produced.
- (c) After examining the data on mRNA half-lives and the amount of protein produced, the researchers hypothesized that each mRNA molecule with modified cap I was translated more frequently than was each mRNA molecule with the normal GTP cap. **Evaluate** their hypothesis by comparing the data in Table 1.
- (d) Introduction of mRNAs into cells allows the cells to produce foreign proteins that they might not normally produce. **Explain** why the production of a foreign protein may be more likely from the introduction of mRNA than DNA into cells.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

STOP

END OF EXAM