



## AP<sup>®</sup> Chemistry 2002 Free-Response Questions

**The materials included in these files are intended for use by AP teachers for course and exam preparation in the classroom; permission for any other use must be sought from the Advanced Placement Program<sup>®</sup>. Teachers may reproduce them, in whole or in part, in limited quantities, for face-to-face teaching purposes but may not mass distribute the materials, electronically or otherwise. These materials and any copies made of them may not be resold, and the copyright notices must be retained as they appear here. This permission does not apply to any third-party copyrights contained herein.**

These materials were produced by Educational Testing Service<sup>®</sup> (ETS<sup>®</sup>), which develops and administers the examinations of the Advanced Placement Program for the College Board. The College Board and Educational Testing Service (ETS) are dedicated to the principle of equal opportunity, and their programs, services, and employment policies are guided by that principle.

The College Board is a national nonprofit membership association dedicated to preparing, inspiring, and connecting students to college and opportunity. Founded in 1900, the association is composed of more than 4,200 schools, colleges, universities, and other educational organizations. Each year, the College Board serves over three million students and their parents, 22,000 high schools, and 3,500 colleges, through major programs and services in college admission, guidance, assessment, financial aid, enrollment, and teaching and learning. Among its best-known programs are the SAT<sup>®</sup>, the PSAT/NMSQT<sup>®</sup>, and the Advanced Placement Program<sup>®</sup> (AP<sup>®</sup>). The College Board is committed to the principles of equity and excellence, and that commitment is embodied in all of its programs, services, activities, and concerns.

Copyright © 2002 by College Entrance Examination Board. All rights reserved. College Board, Advanced Placement Program, AP, SAT, and the acorn logo are registered trademarks of the College Entrance Examination Board. APIEL is a trademark owned by the College Entrance Examination Board. PSAT/NMSQT is a registered trademark jointly owned by the College Entrance Examination Board and the National Merit Scholarship Corporation. Educational Testing Service and ETS are registered trademarks of Educational Testing Service.

INFORMATION IN THE TABLE BELOW AND IN THE TABLES ON PAGES 3-5 MAY BE USEFUL IN ANSWERING THE QUESTIONS IN THIS SECTION OF THE EXAMINATION.

DO NOT DETACH FROM BOOK.

## PERIODIC TABLE OF THE ELEMENTS

1	<b>H</b> 1.0079																	2	<b>He</b> 4.0026
3	4																	9	10
<b>Li</b> 6.941	<b>Be</b> 9.012																	<b>F</b> 19.00	<b>Ne</b> 20.179
11	12																	17	18
<b>Na</b> 22.99	<b>Mg</b> 24.30																	<b>Cl</b> 35.453	<b>Ar</b> 39.948
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36		
<b>K</b> 39.10	<b>Ca</b> 40.08	<b>Sc</b> 44.96	<b>Ti</b> 47.90	<b>V</b> 50.94	<b>Cr</b> 52.00	<b>Mn</b> 54.938	<b>Fe</b> 55.85	<b>Co</b> 58.93	<b>Ni</b> 58.69	<b>Cu</b> 63.55	<b>Zn</b> 65.39	<b>Ga</b> 69.72	<b>Ge</b> 72.59	<b>As</b> 74.92	<b>Se</b> 78.96	<b>Br</b> 79.90	<b>Kr</b> 83.80		
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54		
<b>Rb</b> 85.47	<b>Sr</b> 87.62	<b>Y</b> 88.91	<b>Zr</b> 91.22	<b>Nb</b> 92.91	<b>Mo</b> 95.94	<b>Tc</b> (98)	<b>Ru</b> 101.1	<b>Rh</b> 102.91	<b>Pd</b> 106.42	<b>Ag</b> 107.87	<b>Cd</b> 112.41	<b>In</b> 114.82	<b>Sn</b> 118.71	<b>Sb</b> 121.75	<b>Te</b> 127.60	<b>I</b> 126.91	<b>Xe</b> 131.29		
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86		
<b>Cs</b> 132.91	<b>Ba</b> 137.33	<b>*La</b> 138.91	<b>Hf</b> 178.49	<b>Ta</b> 180.95	<b>W</b> 183.85	<b>Re</b> 186.21	<b>Os</b> 190.2	<b>Ir</b> 192.2	<b>Pt</b> 195.08	<b>Au</b> 196.97	<b>Hg</b> 200.59	<b>Tl</b> 204.38	<b>Pb</b> 207.2	<b>Bi</b> 208.98	<b>Po</b> (209)	<b>At</b> (210)	<b>Rn</b> (222)		
87	88	89	104	105	106	107	108	109	110	111	112	§Not yet named							
<b>Fr</b> (223)	<b>Ra</b> 226.02	<b>†Ac</b> 227.03	<b>Rf</b> (261)	<b>Db</b> (262)	<b>Sg</b> (263)	<b>Bh</b> (262)	<b>Hs</b> (265)	<b>Mt</b> (266)	§ (269)	§ (272)	§ (277)								
*Lanthanide Series																			
58	59	60	61	62	63	64	65	66	67	68	69	70	71						
<b>Ce</b> 140.12	<b>Pr</b> 140.91	<b>Nd</b> 144.24	<b>Pm</b> (145)	<b>Sm</b> 150.4	<b>Eu</b> 151.97	<b>Gd</b> 157.25	<b>Tb</b> 158.93	<b>Dy</b> 162.50	<b>Ho</b> 164.93	<b>Er</b> 167.26	<b>Tm</b> 168.93	<b>Yb</b> 173.04	<b>Lu</b> 174.97						
90	91	92	93	94	95	96	97	98	99	100	101	102	103						
<b>Th</b> 232.04	<b>Pa</b> 231.04	<b>U</b> 238.03	<b>Np</b> 237.05	<b>Pu</b> (244)	<b>Am</b> (243)	<b>Cm</b> (247)	<b>Bk</b> (247)	<b>Cf</b> (251)	<b>Es</b> (252)	<b>Fm</b> (257)	<b>Md</b> (258)	<b>No</b> (259)	<b>Lr</b> (260)						
†Actinide Series																			

GO ON TO THE NEXT PAGE.

## STANDARD REDUCTION POTENTIALS IN AQUEOUS SOLUTION AT 25°C

Half-reaction	$E^\circ(\text{V})$
$\text{F}_2(\text{g}) + 2 e^- \rightarrow 2 \text{F}^-$	2.87
$\text{Co}^{3+} + e^- \rightarrow \text{Co}^{2+}$	1.82
$\text{Au}^{3+} + 3 e^- \rightarrow \text{Au}(\text{s})$	1.50
$\text{Cl}_2(\text{g}) + 2 e^- \rightarrow 2 \text{Cl}^-$	1.36
$\text{O}_2(\text{g}) + 4 \text{H}^+ + 4 e^- \rightarrow 2 \text{H}_2\text{O}(\text{l})$	1.23
$\text{Br}_2(\text{l}) + 2 e^- \rightarrow 2 \text{Br}^-$	1.07
$2 \text{Hg}^{2+} + 2 e^- \rightarrow \text{Hg}_2^{2+}$	0.92
$\text{Hg}^{2+} + 2 e^- \rightarrow \text{Hg}(\text{l})$	0.85
$\text{Ag}^+ + e^- \rightarrow \text{Ag}(\text{s})$	0.80
$\text{Hg}_2^{2+} + 2 e^- \rightarrow 2 \text{Hg}(\text{l})$	0.79
$\text{Fe}^{3+} + e^- \rightarrow \text{Fe}^{2+}$	0.77
$\text{I}_2(\text{s}) + 2 e^- \rightarrow 2 \text{I}^-$	0.53
$\text{Cu}^+ + e^- \rightarrow \text{Cu}(\text{s})$	0.52
$\text{Cu}^{2+} + 2 e^- \rightarrow \text{Cu}(\text{s})$	0.34
$\text{Cu}^{2+} + e^- \rightarrow \text{Cu}^+$	0.15
$\text{Sn}^{4+} + 2 e^- \rightarrow \text{Sn}^{2+}$	0.15
$\text{S}(\text{s}) + 2 \text{H}^+ + 2 e^- \rightarrow \text{H}_2\text{S}(\text{g})$	0.14
$2 \text{H}^+ + 2 e^- \rightarrow \text{H}_2(\text{g})$	0.00
$\text{Pb}^{2+} + 2 e^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+} + 2 e^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+} + 2 e^- \rightarrow \text{Ni}(\text{s})$	-0.25
$\text{Co}^{2+} + 2 e^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Tl}^+ + e^- \rightarrow \text{Tl}(\text{s})$	-0.34
$\text{Cd}^{2+} + 2 e^- \rightarrow \text{Cd}(\text{s})$	-0.40
$\text{Cr}^{3+} + e^- \rightarrow \text{Cr}^{2+}$	-0.41
$\text{Fe}^{2+} + 2 e^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+} + 3 e^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+} + 2 e^- \rightarrow \text{Zn}(\text{s})$	-0.76
$\text{Mn}^{2+} + 2 e^- \rightarrow \text{Mn}(\text{s})$	-1.18
$\text{Al}^{3+} + 3 e^- \rightarrow \text{Al}(\text{s})$	-1.66
$\text{Be}^{2+} + 2 e^- \rightarrow \text{Be}(\text{s})$	-1.70
$\text{Mg}^{2+} + 2 e^- \rightarrow \text{Mg}(\text{s})$	-2.37
$\text{Na}^+ + e^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+} + 2 e^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Sr}^{2+} + 2 e^- \rightarrow \text{Sr}(\text{s})$	-2.89
$\text{Ba}^{2+} + 2 e^- \rightarrow \text{Ba}(\text{s})$	-2.90
$\text{Rb}^+ + e^- \rightarrow \text{Rb}(\text{s})$	-2.92
$\text{K}^+ + e^- \rightarrow \text{K}(\text{s})$	-2.92
$\text{Cs}^+ + e^- \rightarrow \text{Cs}(\text{s})$	-2.92
$\text{Li}^+ + e^- \rightarrow \text{Li}(\text{s})$	-3.05

GO ON TO THE NEXT PAGE.

## ADVANCED PLACEMENT CHEMISTRY EQUATIONS AND CONSTANTS

### ATOMIC STRUCTURE

$$\Delta E = h\nu$$

$$c = \lambda\nu$$

$$\lambda = \frac{h}{m\nu}$$

$$p = m\nu$$

$$E_n = \frac{-2.178 \times 10^{-18}}{n^2} \text{ joule}$$

$E$  = energy

$\nu$  = frequency

$\lambda$  = wavelength

$p$  = momentum

$v$  = velocity

$n$  = principal quantum number

$m$  = mass

### EQUILIBRIUM

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_b = \frac{[\text{OH}^-][\text{HB}^+]}{[\text{B}]}$$

$$K_w = [\text{OH}^-][\text{H}^+] = 1.0 \times 10^{-14} \text{ @ } 25^\circ\text{C}$$

$$= K_a \times K_b$$

$$\text{pH} = -\log [\text{H}^+], \text{pOH} = -\log [\text{OH}^-]$$

$$14 = \text{pH} + \text{pOH}$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

$$\text{pOH} = \text{p}K_b + \log \frac{[\text{HB}^+]}{[\text{B}]}$$

$$\text{p}K_a = -\log K_a, \text{p}K_b = -\log K_b$$

$$K_p = K_c(RT)^{\Delta n},$$

where  $\Delta n$  = moles product gas – moles reactant gas

Speed of light,  $c = 3.0 \times 10^8 \text{ m s}^{-1}$

Planck's constant,  $h = 6.63 \times 10^{-34} \text{ J s}$

Boltzmann's constant,  $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

Avogadro's number =  $6.022 \times 10^{23} \text{ molecules mol}^{-1}$

Electron charge,  $e = -1.602 \times 10^{-19} \text{ coulomb}$

1 electron volt per atom =  $96.5 \text{ kJ mol}^{-1}$

#### Equilibrium Constants

$K_a$  (weak acid)

$K_b$  (weak base)

$K_w$  (water)

$K_p$  (gas pressure)

$K_c$  (molar concentrations)

$S^\circ$  = standard entropy

$H^\circ$  = standard enthalpy

$G^\circ$  = standard free energy

$E^\circ$  = standard reduction potential

$T$  = temperature

$n$  = moles

$m$  = mass

$q$  = heat

$c$  = specific heat capacity

$C_p$  = molar heat capacity at constant pressure

1 faraday  $\mathcal{F}$  = 96,500 coulombs

### THERMOCHEMISTRY

$$\Delta S^\circ = \sum S^\circ \text{ products} - \sum S^\circ \text{ reactants}$$

$$\Delta H^\circ = \sum \Delta H_f^\circ \text{ products} - \sum \Delta H_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \sum \Delta G_f^\circ \text{ products} - \sum \Delta G_f^\circ \text{ reactants}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

$$= -RT \ln K = -2.303 RT \log K$$

$$= -n \mathcal{F} E^\circ$$

$$\Delta G = \Delta G^\circ + RT \ln Q = \Delta G^\circ + 2.303 RT \log Q$$

$$q = mc\Delta T$$

$$C_p = \frac{\Delta H}{\Delta T}$$

**GO ON TO THE NEXT PAGE.**

## GASES, LIQUIDS, AND SOLUTIONS

$$PV = nRT$$

$$\left(P + \frac{n^2a}{V^2}\right)(V - nb) = nRT$$

$$P_A = P_{total} \times X_A, \text{ where } X_A = \frac{\text{moles A}}{\text{total moles}}$$

$$P_{total} = P_A + P_B + P_C + \dots$$

$$n = \frac{m}{M}$$

$$K = ^\circ\text{C} + 273$$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$D = \frac{m}{V}$$

$$u_{rms} = \sqrt{\frac{3kT}{m}} = \sqrt{\frac{3RT}{M}}$$

$$KE \text{ per molecule} = \frac{1}{2}mv^2$$

$$KE \text{ per mole} = \frac{3}{2}RT$$

$$\frac{r_1}{r_2} = \sqrt{\frac{M_2}{M_1}}$$

molarity,  $M$  = moles solute per liter solution

molality = moles solute per kilogram solvent

$$\Delta T_f = iK_f \times \text{molality}$$

$$\Delta T_b = iK_b \times \text{molality}$$

$$\pi = \frac{nRT}{V} i$$

$P$  = pressure

$V$  = volume

$T$  = temperature

$n$  = number of moles

$D$  = density

$m$  = mass

$v$  = velocity

$u_{rms}$  = root-mean-square speed

$KE$  = kinetic energy

$r$  = rate of effusion

$M$  = molar mass

$\pi$  = osmotic pressure

$i$  = van't Hoff factor

$K_f$  = molal freezing-point depression constant

$K_b$  = molal boiling-point elevation constant

$Q$  = reaction quotient

$I$  = current (amperes)

$q$  = charge (coulombs)

$t$  = time (seconds)

$E^\circ$  = standard reduction potential

$K$  = equilibrium constant

$$\begin{aligned} \text{Gas constant, } R &= 8.31 \text{ J mol}^{-1} \text{ K}^{-1} \\ &= 0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1} \\ &= 8.31 \text{ volt coulomb mol}^{-1} \text{ K}^{-1} \end{aligned}$$

$$\text{Boltzmann's constant, } k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$K_f \text{ for H}_2\text{O} = 1.86 \text{ K kg mol}^{-1}$$

$$K_b \text{ for H}_2\text{O} = 0.512 \text{ K kg mol}^{-1}$$

$$1 \text{ atm} = 760 \text{ mm Hg}$$

$$= 760 \text{ torr}$$

$$\text{STP} = 0.000^\circ\text{C and } 1.000 \text{ atm}$$

$$\text{Faraday's constant, } \mathcal{F} = 96,500 \text{ coulombs per mole of electrons}$$

## OXIDATION-REDUCTION; ELECTROCHEMISTRY

$$Q = \frac{[C]^c [D]^d}{[A]^a [B]^b}, \text{ where } a \text{ A} + b \text{ B} \rightarrow c \text{ C} + d \text{ D}$$

$$I = \frac{q}{t}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ - \frac{RT}{n\mathcal{F}} \ln Q = E_{\text{cell}}^\circ - \frac{0.0592}{n} \log Q @ 25^\circ\text{C}$$

$$\log K = \frac{nE^\circ}{0.0592}$$

GO ON TO THE NEXT PAGE.

# 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

## CHEMISTRY

### Section II

(Total time—90 minutes)

#### Part A


Time—40 minutes

**YOU MAY USE YOUR CALCULATOR FOR PART A.**

CLEARLY SHOW THE METHOD USED AND THE STEPS INVOLVED IN ARRIVING AT YOUR ANSWERS. It is to your advantage to do this, since you may obtain partial credit if you do and you will receive little or no credit if you do not. Attention should be paid to significant figures.

Be sure to write all your answers to the questions on the lined pages following each question in the booklet with the pink cover. Do NOT write your answers on the green insert.

Answer Question 1 below. The Section II score weighting for this question is 20 percent.




- Hypobromous acid, HOBr, is a weak acid that dissociates in water, as represented by the equation above.
  - Calculate the value of  $[\text{H}^+]$  in an HOBr solution that has a pH of 4.95.
  - Write the equilibrium constant expression for the ionization of HOBr in water, then calculate the concentration of HOBr(aq) in an HOBr solution that has  $[\text{H}^+]$  equal to  $1.8 \times 10^{-5} M$ .
  - A solution of Ba(OH)<sub>2</sub> is titrated into a solution of HOBr.
    - Calculate the volume of 0.115 M Ba(OH)<sub>2</sub>(aq) needed to reach the equivalence point when titrated into a 65.0 mL sample of 0.146 M HOBr(aq).
    - Indicate whether the pH at the equivalence point is less than 7, equal to 7, or greater than 7. Explain.
  - Calculate the number of moles of NaOBr(s) that would have to be added to 125 mL of 0.160 M HOBr to produce a buffer solution with  $[\text{H}^+] = 5.00 \times 10^{-9} M$ . Assume that volume change is negligible.
  - HOBr is a weaker acid than HBrO<sub>3</sub>. Account for this fact in terms of molecular structure.

## 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

Answer EITHER Question 2 below OR Question 3 printed on page 8. Only one of these two questions will be graded. If you start both questions, be sure to cross out the question you do not want graded. The Section II score weighting for the question you choose is 20 percent.


2. Answer parts (a) through (e) below, which relate to reactions involving silver ion,  $\text{Ag}^+$ .

The reaction between silver ion and solid zinc is represented by the following equation.



- (a) A 1.50 g sample of Zn is combined with 250. mL of 0.110 M  $\text{AgNO}_3$  at 25°C.
- Identify the limiting reactant. Show calculations to support your answer.
  - On the basis of the limiting reactant that you identified in part (i), determine the value of  $[\text{Zn}^{2+}]$  after the reaction is complete. Assume that volume change is negligible.
- (b) Determine the value of the standard potential,  $E^\circ$ , for a galvanic cell based on the reaction between  $\text{AgNO}_3(aq)$  and solid Zn at 25°C.


Another galvanic cell is based on the reaction between  $\text{Ag}^+(aq)$  and  $\text{Cu}(s)$ , represented by the equation below. At 25°C, the standard potential,  $E^\circ$ , for the cell is 0.46 V.



- (c) Determine the value of the standard free-energy change,  $\Delta G^\circ$ , for the reaction between  $\text{Ag}^+(aq)$  and  $\text{Cu}(s)$  at 25°C.
- (d) The cell is constructed so that  $[\text{Cu}^{2+}]$  is 0.045 M and  $[\text{Ag}^+]$  is 0.010 M. Calculate the value of the potential,  $E$ , for the cell.
- (e) Under the conditions specified in part (d), is the reaction in the cell spontaneous? Justify your answer.

## 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

3. Consider the hydrocarbon pentane,  $C_5H_{12}$  (molar mass 72.15 g).
- Write the balanced equation for the combustion of pentane to yield carbon dioxide and water.
  - What volume of dry carbon dioxide, measured at  $25^\circ\text{C}$  and 785 mm Hg, will result from the complete combustion of 2.50 g of pentane?
  - The complete combustion of 5.00 g of pentane releases 243 kJ of heat. On the basis of this information, calculate the value of  $\Delta H$  for the complete combustion of one mole of pentane.
  - Under identical conditions, a sample of an unknown gas effuses into a vacuum at twice the rate that a sample of pentane gas effuses. Calculate the molar mass of the unknown gas.
  - The structural formula of one isomer of pentane is shown below. Draw the structural formulas for the other two isomers of pentane. Be sure to include all atoms of hydrogen and carbon in your structures.





# 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

## CHEMISTRY

### Part B


Time—50 minutes

**NO CALCULATORS MAY BE USED FOR PART B.**

Answer Question 4 below. The Section II score weighting for this question is 15 percent.

4. Write the formulas to show the reactants and the products for any FIVE of the laboratory situations described below. Answers to more than five choices will not be graded. In all cases, a reaction occurs. Assume that solutions are aqueous unless otherwise indicated. Represent substances in solution as ions if the substances are extensively ionized. Omit formulas for any ions or molecules that are unchanged by the reaction. You need not balance the equations.

Example: A strip of magnesium is added to a solution of silver nitrate.




- (a) A solution of sodium iodide is added to a solution of lead(II) acetate.
- (b) Pure solid phosphorus (white form) is burned in air.
- (c) Solid cesium oxide is added to water.
- (d) Excess concentrated hydrochloric acid is added to a 1.0 M solution of cobalt(II) chloride.
- (e) Solid sodium hydrogen carbonate (sodium bicarbonate) is strongly heated.
- (f) An excess of hydrochloric acid is added to solid zinc sulfide.
- (g) Acidified solutions of potassium permanganate and iron(II) nitrate are mixed together.
- (h) A solution of potassium hydroxide is added to solid ammonium chloride.

## 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

Your responses to the rest of the questions in this part of the examination will be graded on the basis of the accuracy and relevance of the information cited. Explanations should be clear and well organized. Examples and equations may be included in your responses where appropriate. Specific answers are preferable to broad, diffuse responses.

Answer BOTH Question 5 below AND Question 6 printed on page 11. Both of these questions will be graded. The Section II score weighting for these questions is 30 percent (15 percent each).



5. A student is asked to determine the molar enthalpy of neutralization,  $\Delta H_{\text{neut}}$ , for the reaction represented above. The student combines equal volumes of 1.0 M HCl and 1.0 M NaOH in an open polystyrene cup calorimeter. The heat released by the reaction is determined by using the equation  $q = mc\Delta T$ .

Assume the following.


- Both solutions are at the same temperature before they are combined.
  - The densities of all the solutions are the same as that of water.
  - Any heat lost to the calorimeter or to the air is negligible.
  - The specific heat capacity of the combined solutions is the same as that of water.
- (a) Give appropriate units for each of the terms in the equation  $q = mc\Delta T$ .
- (b) List the measurements that must be made in order to obtain the value of  $q$ .
- (c) Explain how to calculate each of the following.
- (i) The number of moles of water formed during the experiment
  - (ii) The value of the molar enthalpy of neutralization,  $\Delta H_{\text{neut}}$ , for the reaction between HCl(aq) and NaOH(aq)
- (d) The student repeats the experiment with the same equal volumes as before, but this time uses 2.0 M HCl and 2.0 M NaOH.
- (i) Indicate whether the value of  $q$  increases, decreases, or stays the same when compared to the first experiment. Justify your prediction.
  - (ii) Indicate whether the value of the molar enthalpy of neutralization,  $\Delta H_{\text{neut}}$ , increases, decreases, or stays the same when compared to the first experiment. Justify your prediction.
- (e) Suppose that a significant amount of heat were lost to the air during the experiment. What effect would this have on the calculated value of the molar enthalpy of neutralization,  $\Delta H_{\text{neut}}$ ? Justify your answer.

## 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS

6. Use the principles of atomic structure and/or chemical bonding to explain each of the following. In each part, your answer must include references to both substances.
- (a) The atomic radius of Li is larger than that of Be.
  - (b) The second ionization energy of K is greater than the second ionization energy of Ca.
  - (c) The carbon-to-carbon bond energy in C<sub>2</sub>H<sub>4</sub> is greater than it is in C<sub>2</sub>H<sub>6</sub>.
  - (d) The boiling point of Cl<sub>2</sub> is lower than the boiling point of Br<sub>2</sub>.


Answer EITHER Question 7 below OR Question 8 printed on page 12. Only one of these two questions will be graded. If you start both questions, be sure to cross out the question you do not want graded. The Section II score weighting for the question you choose is 15 percent.

7. An environmental concern is the depletion of O<sub>3</sub> in Earth's upper atmosphere, where O<sub>3</sub> is normally in equilibrium with O<sub>2</sub> and O. A proposed mechanism for the depletion of O<sub>3</sub> in the upper atmosphere is shown below.



- (a) Write a balanced equation for the overall reaction represented by Step I and Step II above.
- (b) Clearly identify the catalyst in the mechanism above. Justify your answer.
- (c) Clearly identify the intermediate in the mechanism above. Justify your answer.
- (d) If the rate law for the overall reaction is found to be  $rate = k[\text{O}_3][\text{Cl}]$ , determine the following.
  - (i) The overall order of the reaction
  - (ii) Appropriate units for the rate constant,  $k$
  - (iii) The rate-determining step of the reaction, along with justification for your answer

## 2002 AP<sup>®</sup> CHEMISTRY FREE-RESPONSE QUESTIONS




8. Carbon (graphite), carbon dioxide, and carbon monoxide form an equilibrium mixture, as represented by the equation above.

- (a) Predict the sign for the change in entropy,  $\Delta S$ , for the reaction. Justify your prediction.
- (b) In the table below are data that show the percent of CO in the equilibrium mixture at two different temperatures. Predict the sign for the change in enthalpy,  $\Delta H$ , for the reaction. Justify your prediction.

Temperature	% CO
700°C	60
850°C	94

- (c) Appropriately complete the potential energy diagram for the reaction by finishing the curve on the graph below. Also, clearly indicate  $\Delta H$  for the reaction on the graph.



- (d) If the initial amount of C(s) were doubled, what would be the effect on the percent of CO in the equilibrium mixture? Justify your answer.

**END OF EXAMINATION**